ewsdendst.com

BREAKING NEWS

LATEST HEADLINES

War of words over operating systems' safety

Exotic black holes spawn new universal law Remains of ancient Egyptian seafaring ships discovered

Public safety issues inform shuttle landing plans Unapproved GM corn found in US food chain Wrinkles could be less than skin deep

Roque weeds defy rules of genetics

First membrane-free alkaline fuel cell built ALL LATEST NEWS

PRINT EDITION

Subscribe



- Current issue
- Archive

Premium Content

JOBS

JOB OF THE WEEK

🕑 Idenix Director Pre-Clinical Studies / Toxicology

Postdoctoral Research Fellow

City of Hope Duarte, US California Postdoctoral Fellow or

Research Associate Position 1 4 1

Case Western Reserve University School of Medicine Cleveland, US Ohio

Production Specialist (Southern Blot) Ingenios Targeting Laboratory Stony Brook, US New York

SUBSCRIPTIONS

Recount slashes number of human genes

18:00 20 October 2004 NewScientist.com news service Andy Coghlan

Humans have just 20,000 to 25,000 genes - well down on previous estimates of 27,000 to 40,000, says the latest analysis of the gene-containing portion of the human genome.

And a separate study has found detailed flaws in "shotgun" sequencing, the more rapid of the two methods used to sequence genomes.

The latest gene count reveals that researchers overestimated the number of genes lurking in heavily-duplicated regions of the human genome, which are extremely tricky to sequence because they are repeated DNA sequences.

"Finding the genes is hard, even in a finished sequence," says Bob Waterston of the University of Washington in Seattle, US, and a lead author of the latest genome analysis.

"But the main point is that we have the genome correct now," he says. "We've got through the many hard parts of the genome, and these included recently duplicated segments which were impossible to study in the earlier drafts.

Evolution and disease

By analysing these "difficult" duplicated regions, Waterston and his colleagues in the International Human Genome Sequencing Consortium discovered that 1183 of the genes had recently been acquired through duplication and the evolution of pre-existing genes.

"These segments are rapidly evolving and appear to be particularly prominent in primate and great ape genomes," says Waterston. "We can at last begin to learn about these and find out how they've contributed to evolution and disease."

A second study has identified shortcomings in shotgun sequencing, the quick fire method of sequencing invented by Craig Venter, founder of Celera Genomics in Rockville, Maryland, US.

Instead of painstakingly amplifying and sequencing each segment of human DNA in cloned bacteria - the traditional method of sequencing - Venter found a way to sequence an entire genome in one go. He did this by smashing it into fragments then reassembling it into the correct order with powerful computer algorithms.

Venter's accelerated method of sequencing precipitated a race between his team at Celera and an international team of publicly funded scientists. But the teams drew in February 2001.

Method of choice

However, the latest analysis - a comparison of Venter's original shotgun sequence with a recent

Related Articles

Bumper compendium of human genes released 20 April 2004

Final human genome sequence released

14 April 2003 First analysis of human DNA reveals fewer

genes than expected 12 February 2001

Search New Scientist Contact us

Web Links

International Human Genome Sequencing Consortium

Robert Waterston, University of Washington J Craig Venter Institute Nature









Make your subscription last longer







username: password :

password?

÷.

SUBSCRIBER LOGIN

Forgotten your

Subscribe now

Athens login

Institutional Subscribers

Log in

SEARCH

Free E-Zine Subscribe to Magazine **Customer Service**



HOME | NEWS | EXPLORE BY SUBJECT | BACK PAGE | SUBSCRIBE | SEARCH | ARCHIVE | RSS | JOBS

The World's No.1 Science & Technology News Service

